

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.


Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 :

C09B 47/00, A61K 31/40

A1

(11) International Publication Number:

WO 94/04614

(43) International Publication Date:

3 March 1994 (03.03.94)

(21) International Application Number: PCT/US93/07686

(22) International Filing Date: 16 August 1993 (16.08.93)

(30) Priority data:

07/929,943

14 August 1992 (14.08.92)

US

08/064,468

20 May 1993 (20.05.93)

US

(71) Applicant: THE TRUSTEES OF THE UNIVERSITY OF PENNSYLVANIA [US/US]; Center for Technology Transfer, 3700 Market Street, Suite 300, Philadelphia, PA 19104 (US).

(74) Agents: CALDWELL, John, C. et al.; Woodcock Washburn Kurtz Mackiewicz & Norris, One Liberty Place - 46th Floor, Philadelphia, PA 19103 (US).


(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published

With international search report.

(72) Inventors: THERIEN, Michael, J. ; 2411 D Delancey Place, Philadelphia, PA 19103 (US). DiMAGNO, Stephen, G. ; 738 Pine Street, Apt. A, Philadelphia, PA 19106 (US).

(54) Title: METAL-MEDIATED CROSS-COUPLING WITH RING-METALATED PORPHYRINS



(57) Abstract

Porphyrins substituted with, for example, vinyl and acetylene groups are provided, along with...

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FR	France	MR	Mauritania
AU	Australia	GA	Gabon	MW	Malawi
BB	Barbados	GB	United Kingdom	NE	Niger
BE	Belgium	GN	Guinea	NL	Netherlands
BF	Burkina Faso	GR	Greece	NO	Norway
BG	Bulgaria	HU	Hungary	NZ	New Zealand
BJ	Benin	IE	Ireland	PL	Poland
BR	Brazil	IT	Italy	PT	Portugal
BY	Belarus	JP	Japan	RO	Romania
CA	Canada	KP	Democratic People's Republic of Korea	RU	Russian Federation
CF	Central African Republic	KR	Republic of Korea	SD	Sudan
CG	Congo	KZ	Kazakhstan	SE	Sweden
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovak Republic
CM	Cameroon	LU	Luxembourg	SN	Senegal
CN	China	LV	Latvia	TD	Chad
CS	Czechoslovakia	MC	Monaco	TC	Togo
CZ	Czech Republic	MG	Madagascar	UA	Ukraine
DE	Germany	ML	Mali	US	United States of America
DK	Denmark	MN	Mongolia	UZ	Uzbekistan
ES	Spain			VN	Viet Nam
FI	Finland				

METAL-MEDIATED CROSS-COUPLING
WITH RING-METALATED PORPHYRINS

RELATED APPLICATION

This patent application is a continuation-in-part
5 of United States Application Serial No. 929,943, filed August
14, 1992. The entire contents of that patent application are
incorporated herein by reference.

FIELD OF THE INVENTION

10 This invention relates to substituted porphyrins and
to techniques and intermediates useful in preparing
substituted porphyrins. The invention also relates to
polymers and macromolecules prepared from the porphyrins, and
to new and improved uses for the porphyrins and their
15 polymers.

BACKGROUND OF THE INVENTION

Porphyrins are derivatives of porphine, a conjugated
cyclic structure of four pyrrole rings linked through their
2- and 5-positions by methine bridges. Porphyrins can be
20 covalently attached to other molecules. The electronic
features of the porphyrin ring system can be altered by the
attachment of one or more substituents. The term "porphyrin"
includes derivatives wherein a metal atom is inserted into the
ring system, as well as molecular systems in which ligands are
25 attached to the metal. The substituents, as well as the
overall porphyrin structure, can be neutral, positively
charged, or negatively charged.

-2-

Numerous porphyrins have been isolated from natural sources. Notable porphyrin-containing natural products include hemoglobin, the chlorophylls, and vitamin B₁₂. Also, many porphyrins have been synthesized in the laboratory, typically through condensation of suitably substituted pyrroles and aldehydes. However, reactions of this type generally proceed in low yield, and cannot be used to produce many types of substituted porphyrins.

Accordingly, there exists a need in the art for synthetic methods capable of producing a greater variety of porphyrins than presently available.

OBJECTS OF THE INVENTION

It is one object of the present invention to provide improved methods for synthesizing substituted porphyrins.

It is another object of the invention to provide novel substituted porphyrins.

It is yet another object to provide novel porphyrin-containing compounds.


It is a further object of the invention to provide polymers containing linked porphyrin units.

It is still another object to provide new applications for substituted porphyrins and porphyrin-containing compounds.


SUMMARY OF THE INVENTION

These and other objects are satisfied by the present invention, which provides porphyrins having formula (1), (2), or (3):


-3-



(1)



(2)



(3)

wherein M and M' are metal atoms and R_{A1} - R_{A4} and R_{B1} - R_{B8} are, independently, H or chemical functional groups that can bear a negative charge prior to attachment to a porphyrin compound. In certain preferred embodiments, at least one of R_{A1} - R_{A4} has formula $CH=CH_2$, or at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C(R_C)=C(R_D)(R_E)$, provided that at least one of R_C , R_D , and R_E is not H, where R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 4 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl or porphyrinato; M is a transition metal, a lanthanide, actinide, rare earth or alkaline metal. R_C , R_D , and R_E also can include peptides, nucleosides, and/or saccharides.

-4-

In other preferred embodiments, at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C\equiv C(R_D)$. In further preferred embodiments, at least one of R_{A1} - R_{A4} is haloalkyl having from 1 to about 20 carbon atoms. In further preferred embodiments,
 5 at least one of R_{B1} - R_{B8} is haloalkyl having 2 to about 20 carbon atoms or at least five of R_{B1} - R_{B8} are haloalkyl having from 1 to about 20 carbon atoms or haloaryl having from about 4 to about 20 carbon atoms. In further preferred embodiments, at least one of R_{B1} - R_{B8} is haloaryl or haloheteroaryl having
 10 about 4 to about 20 carbon atoms. In still further preferred embodiments, at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} includes an amino acid, peptide, nucleoside, or saccharide.

The present invention also provides processes and intermediates for preparing substituted porphyrins. In
 15 certain embodiments, the processes comprise providing a porphyrin compound having formula (1), (2), or (3) wherein at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} is a halogen and contacting the porphyrin compound with a complex having formula $Y(L)_2$ wherein Y is a metal and L is a ligand. This produces a first
 20 reaction product, which is contacted with an organometallic compound having general formula $T(R_L)_z(R_O)_y(X_B)_w$, $T(R_O)_y(X_B)$ or $T(R_O)_y$ where T is a metal; X_B is a halogen; R_L is cyclopentadienyl or aryl having about 6 to about 20 carbon atoms; R_O is alkyl having 1 to about 10 carbon atoms, alkenyl
 25 or alkynyl having 2 to about 10 carbon atoms, aryl having about 6 to about 20 carbon atoms; z and w are greater than or equal to 0; and y is at least 1. This contacting produces a second reaction product which, through reductive elimination, yields a third reaction product that contains a porphyrin
 30 substituted with R_O .

In further embodiments, the processes comprise providing a porphyrin compound having formula (1), (2), or (3) wherein at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} is a halogen and contacting the porphyrin compound with an activated metal Z.
 35 This produces a ring-metalated porphyrin wherein metal Z has inserted into the porphyrin/halogen covalent bond. Alternatively, ring metalation of the halogenated porphyrin

-5-

can be carried out with a catalytic amount of $Y(L)_2$ and an alkyltin reagent having formula $(R)_3Sn-Sn(R)_3$, where R is alkyl having 1 to about 10 carbon atoms, to yield trialkyltin-substituted porphyrin. Either metalated porphyrin can be

5 contacted with a compound having formula R_p-R_z , where R_p is cyclopentadienyl or aryl having about 6 to about 20 carbon atoms, alkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, or porphyrinato and R_z is a leaving group such as a halogen, a tosylate, or a triflate.

10 Such contacting should be performed in the presence of a catalyst having formula $Y(L)_2$, as defined above. This produces a porphyrin substituted with R_p . Alternatively, the first reaction product can be contacted with an organic or organometallic quenching reagent.

15 In another aspect, the invention provides polymers comprising linked porphyrin units. In certain embodiments, porphyrin units having formula (1), (2), or (3) share covalent bonds. In other embodiments, at least one of $R_{A1}-R_{A4}$ or $R_{B1}-R_{B8}$ function as linking groups. In these embodiments, at least

20 a portion of a linking group can have formula $[C(R_C)=C(R_D)(R_E)]_x$, $[C\equiv C(R_D)]_x$, $[CH_2(R_C)-CH(R_D)(R_E)]_x$ or $[CH=CH(R_D)]_x$ where x is at least 1. The remaining of $R_{A1}-R_{A4}$ and $R_{B1}-R_{B8}$ can be H, halogen, alkyl or heteroalkyl having 1 to about 20 carbon atoms or aryl or heteroaryl having 4 to about

25 20 carbon atoms, $C(R_C)=C(R_D)(R_E)$, $C\equiv C(R_D)$, or a chemical functional group that includes a peptide, nucleoside, and/or saccharide. In other preferred embodiments, the linking group is cycloalkyl or aryl having about 6 to about 22 carbon atoms.

The invention also provides processes for preparing

30 porphyrin-containing polymers. In certain embodiments, the processes comprise providing at least two compounds that, independently, have formula (1), (2), or (3) wherein at least one of $R_{A1}-R_{A4}$ or $R_{B1}-R_{B8}$ in each of the compounds contains an olefinic carbon-carbon double bond or a chemical functional

35 group reactive therewith. In other embodiments, at least one of $R_{A1}-R_{A4}$ or $R_{B1}-R_{B8}$ in each of the compounds contains a carbon-carbon triple bond or a chemical functional group reactive

-6-

therewith. The compounds are then contacted for a time and under reaction conditions effective to form covalent bonds through the carbon-carbon double and/or triple bonds.

The porphyrins and porphyrin-containing polymers of the invention can be used, for example, as dyes, catalysts, contrast agents, antitumor agents, antiviral agents, and in chemical sensors and electrooptical devices.

BRIEF DESCRIPTION OF THE FIGURES

The numerous objects and advantages of the present invention may be better understood by those skilled in the art by reference to the accompanying figures, in which:

Figure 1 shows a linear polymer of the invention wherein R_{F1} is a covalent bond or a divalent functional group.

Figure 2 shows a linear polymer of the invention having diacetylenic linking groups.


Figure 3 shows a linear polymer of the invention having monoacetylenic and phenoxy linking groups, wherein R' is alkyl having from 1 to about 20 carbon atoms.

Figure 4 shows a cross-linked polymer of the invention, wherein n , m , v , x , and y are at least 1.


DETAILED DESCRIPTION OF THE INVENTION

It has been found in accordance with the present invention that a wide variety of novel porphyrins can be prepared through metal-mediated cross coupling of a halogenated porphyrin core and a suitable organometallic moiety. In general, the resulting porphyrins have formula (1), (2), or (3):


-7-



(1)



(2)



(3)

wherein M and M' are metal atoms and R_{A1} - R_{A4} and R_{B1} - R_{B8} are, independently, H or a variety of chemical functional groups that can bear a negative charge prior to attachment to a porphyrin compound. M preferably is a lanthanide or actinide
 5 or a metal such as Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Y, Zr, Nb, Mo, Ru, Rh, Pd, Ag, La, Hf, Ta, W, Re, Os, Ir, Pt, Cd, Hg, Li or Au. More preferably, M is a metal having a full valence shell, even more preferably Zn or Mg. M' can be a metal such as Li, Na, K, Rb, or Cs, preferably Li.

10 R_{A1} - R_{A4} and R_{B1} - R_{B8} can be virtually any chemical functional group or covalent assemblage of functional groups which, prior to attachment to a porphyrin compound, can bear a negative charge. In certain embodiments wherein R_{A1} - R_{A4} and R_{B1} - R_{B8} are appended to a porphyrin compound using the methods
 15 of the invention, these groups should be capable of bearing a carbon-centered negative charge (i.e., exist as a

-8-

carbocentric anion in solution or otherwise). Preferably, R_{A1} - R_{A4} and R_{B1} - R_{B8} are primary or secondary alkyl, alkenyl, haloalkenyl, alkynyl, haloalkynyl, aryl or heteroaryl. R_{A1} - R_{A4} and R_{B1} - R_{B8} should contain functionality that can withstand the
5 reaction conditions associated with metal-mediated cross coupling. Those skilled in the art will recognize that chemical protecting groups can be attached to any sensitive functionality found within R_{A1} - R_{A4} and R_{B1} - R_{B8} and then removed after the coupling reactions have been completed.

10 Compounds having formulas (1)-(3) preferably bear 1, 2, 4, 8, or 12 substituents (i.e., 1, 2, 4, 8, or 12 of R_{A1} - R_{A4} and R_{B1} - R_{B8} are not H). ~~More preferably, these compounds~~
bear 2 or 4 substituents that contain alkenyl or alkynyl functionality.

15 In certain embodiments, at least one of R_{A1} - R_{A4} has formula $CH=CH_2$. In other embodiments, at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C(R_C)=C(R_D)(R_E)$, provided that at least one of R_C , R_D , and R_E is not H, or at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C\equiv C(R_D)$. In further preferred embodiments,
20 at least one of R_{A1} - R_{A4} is haloalkyl having from 1 to about 20 carbon atoms. In further preferred embodiments, at least one of R_{B1} - R_{B8} is haloalkyl having 2 to about 20 carbon atoms or at least five of R_{B1} - R_{B8} are haloalkyl having from 1 to about 20 carbon atoms. In further preferred embodiments, at least one
25 of R_{B1} - R_{B8} is haloaryl or haloheteroaryl having about 4 to about 20 carbon atoms. Preferred halogenated moieties are fully halogenated (e.g., CF_3). In still further embodiments, at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} is a single amino acid or a terminal amino acid of a polypeptide.

30 R_C , R_D , and R_E can be any of a wide variety of chemical functional groups. Preferably, R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 4 to about 20 carbon atoms, alkenyl or heteroalkenyl
35 having from 2 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 2 to about 20 carbon atoms, trialkylsilyl, or porphyrinato. The terms heteroalkyl and

-9-

heteroaryl are intended to denote moieties wherein a heteroatom is inserted into the carbon backbone of an alkyl or aryl structure (e.g., ether, thioether, and pyridinyl groups). Representative heteroatoms include N, O, S, Se, and Te. The terms alkyl, aryl, alkenyl, and alkynyl are intended to include moieties substituted with, for example, halogens or nitro, acid, alkyl, or ester groups. Preferred alkyl and aryl groups have from 1 to about 10 carbon atoms and about 6 to about 14 carbon atoms, respectively. Preferred alkenyl and alkynyl groups have from 2 to about 10 carbon atoms. R_C , R_D , and R_E preferably are alkyl, aryl, or trialkylsilyl.

R_C , R_D , and/or R_E also can be chemical functional groups that include at least one peptide, nucleoside, and/or saccharide. For example, R_C , R_D , and R_E can include a polymethylene chain connecting a DNA- or RNA-cleaving compound having formula (1), (2), or, in the absence of water, (3) with an oligonucleotide. (see, e.g., Stein, et al., *Cancer Research* 1988, 48, 2659). As will be recognized, peptides are compounds comprising two or more amino acids covalently bound through an amide linkage (e.g., glycylalanine), nucleosides are glycosides comprising covalently bound pentoses and heterocyclic bases (e.g., cytidine), and saccharides are hemiacetal forms of polyhydroxy aldehydes and ketones (e.g., sucrose). Each of these terms is intended to include both naturally occurring and non-naturally occurring moieties.

In preferred embodiments, each of R_{A1} - R_{A4} is alkyl or aryl and either R_{B5} , R_{B5} and R_{B1} , or R_{B5} and R_{B2} have formula $C\equiv C(R_D)$. In other preferred embodiments, each of R_{A1} - R_{A4} is alkyl or aryl and each of R_{B1} - R_{B8} have formula $C\equiv C(R_D)$. In still other preferred embodiments, R_{A1} has formula $C\equiv C(R_D)$, each of R_{A2} - R_{A4} are alkyl or aryl, at least one of R_{B1} - R_{B8} is H, alkyl or aryl, and R_D is H, F, Cl, Br, I, alkyl, aryl, alkenyl, alkynyl, trialkylsilyl, or porphyrinato. In other preferred embodiments, R_{A1} and R_{A3} have formula $C\equiv C(R_D)$, R_{A2} and R_{A4} are alkyl, aryl or a halogen, at least one of R_{B1} - R_{B8} is H, alkyl, aryl or a halogen, and R_D is H, F, Cl, Br, I, alkyl, aryl, alkenyl, alkynyl, trialkylsilyl, or porphyrinato; more

-10-

preferably R_D is 3,4,5-trialkyl-substituted phenyl or 3,4,5-trialkyl-oxy-substituted phenyl with the alkyl and alkoxy substituents having about 7 to about 20 carbon atoms. In still other preferred embodiments, R_{A1} - R_{A4} have formula $C\equiv C(R_D)$, at least one of R_{B1} - R_{B8} is H, alkyl, aryl or a halogen, and R_D is H, F, Cl, Br, I, alkyl, aryl, alkenyl, alkynyl, trialkylsilyl, or porphyrinato, more preferably R_D is H, 3,4,5-trialkyl-substituted phenyl, or 3,4,5-trialkyl-oxy-substituted phenyl with the alkyl and alkoxy substituents having about 7 to about 20 carbon atoms.

In certain embodiments, the porphyrins of the invention are prepared by metal-mediated cross-coupling of a halogenated, preferably brominated, chlorinated, or iodinated, porphyrin with an organometallic moiety. In other embodiments, the porphyrins of the invention are prepared by metal-mediated cross coupling of a metalated porphyrin with an organic moiety bearing a suitable leaving group. As will be recognized, the latter methods provide a means for attaching chiral substituents to a porphyrin core. Both metalated porphyrins (e.g., formulas (2) and (3)) and non-metalated porphyrins (e.g., formula (1)) can be used in the processes of the invention; the cross-coupling products can be metalated and de-metalated, as desired. The principles and techniques relating to metal-mediated cross coupling are well known to those skilled in the art to consist of three general steps: (1) oxidative addition, (2) transmetalation, and (3) reductive elimination. (See, e.g., Collman, et al., Principles and Applications of Organotransition Metal Chemistry, University Science Books, 1987, Mill Valley, CA; Kumada, Pure & Appl. Chem., 1980, 52, 669; Hayashi, et al., Tetrahedron Letters, 1979, 21, 1871.)

In accordance with certain embodiments of the invention, a halogenated porphyrin, P_N-X_A , is contacted with a catalyst having formula $Y(L)_2$, where Y is a metal such as, for example, Pd, Ni, Pt, Ru, or Cu and L is a ligand appropriate for that metal. When Y is Pd or Ni, L should be a phosphorous-, nitrogen-, arsenic-, or antimony-containing

-11-

Lewis base such as an alkyl- or arylphosphine, an alkyl or arylarsenine, an alkyl- or arylstibene, a nitrogen-containing heterocycle such as pyridine, or a mixture thereof. The terms alkylphosphine, arylphosphine, alkylarsenine, arylarsenine, alkylstibene, and arylstibene are intended to describe moieties having at least one alkyl or aryl substituent such as, for example, monoalkylphosphines, trialkylphosphines, and dialkylmonoarylpophosphines. Contacting the halogenated porphyrin with the catalyst complex is believed to produce a second compound having formula $P_N-Y(L)_2X_A$, which is contacted with an organometallic compound having formula $T(R_L)_z(R_O)_w$, $T(R_L)_z(R_O)_t(X_B)_v$, $T(R_O)_t(X_B)_v$ or $T(R_O)_t$, where T is a metal such as, for example, Li, Na, K, Rb, Cs, Hg, Sn, Al, B, Si, Zn, Zr, Cd, Cu, or Mg; X_B is a halogen or an alkoxy group; R_L is cyclopentadienyl or aryl having about 6 to about 20 carbon atoms; R_O is alkyl, alkenyl or alkynyl having 1 to about 10 carbon atoms, aryl having about 6 to about 20 carbon atoms such as $C(R_C)=C(R_D)(R_E)$, $C\equiv C(R_D)$, haloalkyl groups or haloaryl groups; z and w are greater than or equal to 0; and t is at least 1. T can be any metal that does not participate in an outer sphere electron transfer reaction with the porphyrin. Representative organometallic compounds are CuR_O , $Zn(R_O)_2$, ZnR_OX_B , $(nBu)_3SnR_O$, and $Cp_2ZrR_OX_B$ (Cp = cyclopentadienyl). This contacting is believed to produce a third compound having formula $P_N-Y(L)_2R_O$ which, through reductive elimination, is transformed into substituted porphyrin P_N-R_O .

In other embodiments of the invention, metalated porphyrins are prepared by contacting halogenated porphyrins, P_N-X_A , with an activated metal Z. Alternatively, ring metalation of the halogenated porphyrin can be carried out with a catalytic amount of $Y(L)_2$ and an alkyltin reagent having formula $(R)_3Sn-Sn(R)_3$, where R is alkyl having 1 to about 10 carbon atoms, to yield trialkyltin-substituted porphyrin. Z can be any metal that does not participate in an outer sphere electron transfer reaction with the porphyrin. Z preferably is Zn, Mg, Ca, Cd, Cu, Sn, Sr, Ba, or Zr. Preferably, the activated metal is prepared by contacting a

-12-

metal reagent having formula $Z(X_c)_c$, where Z is a metal, X_c is a halogen, and c is 1-4, with a reducing agent under conditions effective to reduce the metal reagent. In embodiments wherein the metal reagent is reduced in the presence of the porphyrin compound, the reducing agent should be effective to reduce the metal reagent but ineffective to reduce the porphyrin compound. Activated metals according to the invention can be prepared by, for example, the method of Ebert and Rieke, *J. Org. Chem.* 1984, 49, 5280 and *J. Org. Chem.* 1988, 53, 4482.

The metalated porphyrins can be contacted with compounds having formula R_1-R_2 , where R_1 is cyclopentadienyl or aryl having about 6 to about 20 carbon atoms, alkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, or porphyrinato and R_2 is a leaving group. Leaving groups include but are not limited halogen, alkylsulfonyl, substituted alkylsulfonyl, arylsulfonyl, substituted arylsulfonyl, heterocyclosulfonyl or trichloroacetimidate. Preferred leaving groups include chloro, fluoro, bromo, iodo, p-(2,4-dinitroanilino)benzenesulfonyl, benzenesulfonyl, methylsulfonyl (mesylate), p-methylbenzenesulfonyl (tosylate), p-bromobenzenesulfonyl, trifluoromethylsulfonyl (triflate), trichloroacetimidate, acyloxy, 2,2,2-trifluoroethanesulfonyl, imidazolesulfonyl, and 2,4,6-trichlorophenyl. Contacting should be performed in the presence of a catalyst having formula $Y(L)_2$, where Y is a metal such as, for example, Pd, Ni, Pt, Ru, or Cu and L is a ligand appropriate for that metal. When Y is Pd or Ni, L should be a phosphorous-, nitrogen-, arsenic-, or antimony-containing Lewis base such as an alkyl- or arylphosphine, an alkyl or arylarsenine, an alkyl- or arylstibene, a nitrogen-containing heterocycle such as pyridine, or a mixture thereof.

Ring-metalated porphyrins alternatively can be contacted with organic or organometallic quenching reagents. Representative organic quenching reagents include those commonly used in the well known Grignard reaction, such as, for example, CO_2 , esters, aldehydes, ketones, acid chlorides,

-13-

and amides. (see, e.g., J. March, Advanced Organic Chemistry, John Wiley & Sons, 1985, New York) Organometallic quenching reagents include those disclosed by Azizian, et al., *J. Organomet. Chem.* 1981, 215, 49 and Barbero, et al., *J. Chem. Soc. Chem. Commun.* 1992, 351, such as, for example, trialkyl tin halides and trialkyl borates.

In general, the coupling reactions of the present invention proceed rapidly and in excellent yield, contrary to teaching in the prior art that reactions of this type should not work well with electron-rich systems such as porphyrins. It has been found that not all of the metals known for use in metal-mediated cross coupling reactions can be used to prepare substituted porphyrins. For example, coupling reactions wherein the metal T is lithium or magnesium have been found to proceed, if at all, in very low yield and with destruction of the porphyrin starting material. Also, coupling reactions wherein the metal Z is lithium, potassium, or sodium preferably are avoided.


The use of metal-mediated cross coupling in accordance with the invention can produce monomeric compounds suitable for incorporation into porphyrin-containing homopolymers or copolymers or into macromolecular or supramolecular species containing, for example, one or more peptides, nucleosides, or saccharides. Alternatively, metal-mediated cross-coupling can be used to directly prepare porphyrin-containing homopolymers or copolymers. The polymers according to the invention can contain as few as 2 porphyrin units, but more preferably contain at least 3 porphyrin units, more preferably at least 5 porphyrin units. In certain embodiments, polymers of the invention comprise a plurality of porphyrin units that, independently, have formula (1), (2), or (3) wherein at least one of R_{B1} - R_{B8} or R_{A1} - R_{A4} includes a linking group selected from $[C(R_C)=C(R_D)(R_E)]_x$, $[C\equiv C(R_D)]_x$, $[CH_2(R_C)-CH(R_D)(R_E)]_x$ or $[CH=CH(R_D)]_x$ where x is at least 1. Preferably, the remaining of R_{A1} - R_{A4} are, independently, H, alkyl, alkenyl, alkynyl, or aryl and the remaining of R_{B1} - R_{B8}

-14-

are, independently, H, alkyl, aryl, $C(R_C)=C(R_D)(R_E)$, or $C\equiv C(R_D)$.

In other embodiments, polymers according to the invention comprise a plurality of porphyrin units that, independently, have formula (1), (2), or (3) wherein at least one of $R_{B1}-R_{B8}$ or $R_{A1}-R_{A4}$ is a cycloalkyl, cycloalkenyl, aryl or heteroaryl linking group having about 6 to about 22 carbon atoms.


Those skilled in the art will recognize the wide variety of polymers that can be prepared from the porphyrin-containing compounds of the invention. In certain embodiments, cofacial polymers are formed having, for example, formula (4). (see, e.g., Durand, et al., *J. Am. Chem. Soc.*, 1983, 105, 2710).



(4)

15


In other embodiments, somewhat linear polymer chains are formed wherein a portion of the polymer has general formula $(P_N)_r$, where P_N is a porphyrin unit and r is at least 2. In further embodiments, linear polymer chains have general formula:




where Q_L is a linking group, P_N is a porphyrin unit, and h , 1, and s are independently selected to be at least 1. For example, a portion of such polymers can have formula:

20


-15-



wherein P_{N1} and P_{N2} are independently selected porphyrin units, Q_{L1} and Q_{L2} are independently selected linking groups, and l' , l'' , s' , and s'' are at least 1. These essentially linear
 5 polymer chains can be cross-linked such that a portion of the polymer has general formula:




wherein Q_H is a linking group, and h , u , and v are independently selected to be at least 1. A portion of these
 10 cross-linked polymers can have formula:




wherein P_{N3} is a porphyrin unit, Q_{H1} and Q_{H2} are independently
 selected linking groups, and h' , h'' , u' , and u'' are at least
 1. Thus, cross-linked polymers can have formulas:

15



20



where r' is at least 1.

The polymers of the invention can be formed by
 25 contacting a substituted porphyrin with a second compound containing functionality that is reactive with the functionality contained within the porphyrin. Preferably, the porphyrin contains an olefinic carbon-carbon double bond, a carbon-carbon triple bond or some other reactive
 30 functionality. The contacting should be performed under conditions effective to form a covalent bond between the respective reactive functionalities. Preferably, porphyrin-containing polymers are formed by metal-mediated cross-coupling of, for example, dibrominated porphyrin units. Also,
 35 porphyrin-containing polymers can be synthesized using known terminal alkyne coupling chemistry. (see, e.g., Patai, et al., The Chemistry of Functional Groups, Supplement C, Part 1, pp. 529-534, Wiley, 1983; Cadiot, et al., Acetylenes, pp. 597-647, Marcel Dekker, 1964; and Eglinton, et al., Adv. Org. Chem.,

-16-

1963, 4, 225) As will be recognized, the second compound noted above can be a substituted porphyrin of the invention or some other moiety such as an acrylate monomer. Thus, a wide variety of copolymeric structures can be synthesized with the porphyrins of the invention. Through careful substituent selection the porphyrins of the invention can be incorporated into virtually any polymeric matrix known in the art, including but not limited to polyacetylenes, polyacrylates, polyolefins, polyethers, polyurethanes, polycarbonates, polyanilines, polypyrroles, and polythiophenes. For example, fluorescent porphyrins can be incorporated into such polymers as end-capping groups.

The porphyrins and porphyrin-containing polymers of the invention can be used, for example, as dyes, catalysts, contrast agents, antitumor agents, antiviral agents, liquid crystals, in chemical sensors and in electrooptical and solar energy conversion devices. They also can be incorporated into supramolecular structures. The polymers and supramolecular structures, which anchor porphyrin units in a relatively stable geometry, should improve many of the known uses for porphyrins and even provide a number of new uses, such as in a solid phase system for sterilizing virus-containing solutions. Representative uses are disclosed by, for example, the following patents, which are incorporated herein by reference: U.S. Patent No. 4,895,682 (Ellis, et al.); U.S. Patent No. 4,986,256 (Cohen); U.S. Patent No. 4,668,670 (Rideout, et al.); U.S. Patent No. 3,897,255 (Erickson); U.S. Patent No. 3,899,334 (Erickson); U.S. Patent No. 3,687,863 (Wacher); U.S. Patent No. 4,647,478 (Formanek, et al.); and U.S. Patent No. 4,957,615 (Ushizawa, et al.). Further uses are disclosed are disclosed by, for example, U.K. Patent Application 2,225,963 (Casson, et al.); International Application WO 89/11277 (Dixon, et al.); International Application WO 91/09631 (Matthews, et al.); European Patent Application 85105490.8 (Weishaupt, et al.); European Patent Application 90202953.7 (Terrell, et al.); European Patent Application 89304234.1 (Matsushima, et al.); Lehn, Angew.

-17-

Chem. Int. Ed. Engl., 1988, 27, 89; Wasielewski, Chem. Rev., 1992, 92, 435; Mansury, et al., J. Chem. Soc., Chem. Comm., 1985, 155; Groves, et al., J. Am. Chem. Soc., 1983, 105, 5791; and Giroud-Godquin, et al., Angew. Chem. Int. Ed. Engl., 1991, 30, 375. It is believed that the porphyrins of the invention can be substituted for the porphyrins disclosed in each of the foregoing publications.

Additional objects, advantages, and novel features of this invention will become apparent to those skilled in the art upon examination of the following examples thereof, which are not intended to be limiting.

EXAMPLE 1**5,15-DIPHENYLPORPHYRIN**

A flame-dried 1000 ml flask equipped with a magnetic stirring bar was charged with 2,2-dipyrrylmethane (458 mg, 3.1 mmol), benzaldehyde (315 μ l, 3.1 mmol), and 600 ml of freshly distilled (CaH₂) methylene chloride. The solution was degassed with a stream of dry nitrogen for 10 minutes. Trifluoroacetic acid (150 μ l, 1.95 mmol) was added via syringe, the flask was shielded from light with aluminum foil, and the solution was stirred for two hours at room temperature. The reaction was quenched by the addition of 2,3-dichloro-5,6-dicyano-1,4-benzoquinone (DDQ, 900 mg, 3.96 mmol) and the reaction was stirred for an additional 30 minutes. The reaction mixture was neutralized with 3 ml of triethylamine and poured directly onto a silica gel column (20 x 2 cm) packed in hexane. The product was eluted in 700 ml of solvent. The solvent was evaporated, leaving purple crystals (518 mg., 1.12 mmol, 72.2%). This product was sufficiently pure for further reactions. Vis(CHCl₃): 421 (5.55), 489 (3.63), 521 (4.20), 556 (4.04), 601 (3.71), 658 (3.73).

-18-

EXAMPLE 2

5,15-DIBROMO-10,20-DIPHENYLPORPHYRIN

5,15-Diphenylporphyrin (518 mg, 1.12 mmol) was dissolved in 250 ml of chloroform and cooled to 0°C. Pyridine (0.5 ml) was added to act as an acid scavenger. N-Bromosuccinimide (400 mg, 2.2 mmol) was added directly to the flask and the mixture was followed by thin-layer chromatography (TLC; 50% CH₂Cl₂/hexanes eluant). After 10 minutes the reaction reached completion and was quenched with 20 ml of acetone. The solvents were evaporated and the product was washed with several portions of methanol and pumped dry to yield 587 mg (0.94 mmol, 85%) of reddish-purple solid. The compound was sufficiently pure to use in the next reaction. Vis(CHCl₃): 421 (5.55), 489 (3.63), 521 (4.20), 556 (4.04), 601 (3.71), 658 (3.73).

EXAMPLE 3

5,15-DIBROMO-10,20-DIPHENYLPORPHYRINATO ZINC

5,15-Dibromo-10,20-diphenylporphyrin (587 mg, 0.94 mmol) was suspended in 30 ml DMF containing 500 mg ZnCl₂. The mixture was heated at reflux for 2 hours and poured into distilled water. The precipitated purple solid was filtered through a fine fritted disk and washed with water, methanol, and acetone and dried in vacuo to yield 610 mg (0.89 mmol, 95%) of reddish purple solid. The compound was recrystallized from tetrahydrofuran (THF)/heptane to yield large purple crystals of the title compound (564 mg, 0.82 mmol, 88%). Vis(THF): 428 (5.50), 526 (3.53), 541 (3.66), 564 (4.17), 606 (3.95).

EXAMPLE 4

MESO-SUBSTITUTED PORPHYRINS

General Procedure

In each of the following examples, 5,15-Dibromo-10,20-diphenylporphyrinato zinc (0.1 mmol), and Pd(PPh₃)₄ (0.0025 mmol) were dissolved in 35 ml of distilled, degassed THF in a sealed storage tube with the 1 mmol of the indicated

-19-

organometallic reagent and warmed at 60°C for 48 hours. The reaction was monitored by TLC on withdrawn aliquots. The mixture was quenched with water, extracted with chloroform, dried over CaCl_2 , evaporated and purified by column chromatography.

A. 5,15-Diphenyl-10,20-dimethylporphyrinato zinc

The organometallic reagent was methyl zinc chloride prepared from methyl lithium and anhydrous zinc chloride in THF.

10 The crude solid was dissolved in THF/heptane, poured onto 10 g silica gel and evaporated to dryness. This silica gel was loaded onto a column packed in 50% CH_2Cl_2 /hexane. A single band was eluted (50% CH_2Cl_2 /hexane) to yield pure 5,15-diphenyl-10,20-dimethylporphyrinato zinc (48 mg, 88%). An analytical sample was recrystallized from THF/heptane by slow evaporation under N_2 . ^1H NMR (500 MHz, 3:1 CDCl_3 , D_8 -THF) δ 9.34 (d, 4H, $J=4.6$); 8.71 (d, 4H, $J=4.6$); 8.02 (dd, 4H, $J_1=7.5$, $J_2=1.4$); 7.57 (m, 6H); 4.51 (s, 6H). ^{13}C NMR (125 MHz, 3:1 CDCl_3 , D_8 -THF) δ 150.07(0), 148.88(0), 143.34(0), 134.18(1), 131.42(1), 128.09(1), 126.73(1), 125.88(1), 119.29(0), 113.74(0), 20.81(3). Vis (THF) 424 (5.58), 522 (3.40), 559 (4.12); 605 (3.88).

B. 5,15-Diphenyl-10,20-divinylporphyrinato zinc

The organometallic reagent was tri-n-butylvinyl tin.

25 The crude product was absorbed on silica and loaded onto a column packed in hexane. Elution was carried out with CH_2Cl_2 (0-50%)/hexane. A small quantity of purple material led the main fraction. The main band was evaporated to yield pure 5,15-diphenyl-10,20-divinylporphyrinato zinc (53 mg, 91%). An analytical sample was recrystallized from chloroform. ^1H NMR (500 MHz, CDCl_3) δ 9.52 (d, 4H, $J=4.7$); 9.24(dd, 2H, $J_1=17.3$, $J_2=9.1$); 8.92 (d, 4H, $J=4.7$); 8.19 (dd, 4H, $J_1=6.8$, $J_2=2.0$); 7.75 (m, 6H); 6.48 (dd, 2H, $J_1=11.0$, $J_2=1.9$); 6.05 (dd, 2H, $J_1=17.3$, $J_2=2.0$). ^{13}C NMR (125 MHz, CDCl_3) δ 163.40(1), 149.90(0), 149.21(0), 142.83(0), 137.97(0), 134.40(1), 132.10(1), 130.39(1), 127.50(1), 126.73(2), 126.57(1), 121.05(0).

-20-

C. 5,15-Bis(2,5-dimethoxyphenyl)-10,20-diphenylporphyrinato zinc

The organometallic reagent was 2,5-dimethoxyphenyl lithium, prepared from 1,4-dimethoxybenzene and *t*-butyl lithium in ether at -78°C. The organolithium reagent was added to a solution of ZnCl₂ in THF to yield the organozinc chloride reagent. This reagent was used immediately.

At the completion of the reaction two highly fluorescent spots were visible by TLC. The crude product was chromatographed on silica using CHCl₃ as eluant. The first band off the column proved to be the C_{2h} isomer of 5,15-bis(2,5-dimethoxyphenyl)-10,20-diphenylporphyrinato zinc.

This band was evaporated leaving 33 mg (42%) of pure product. An analytical sample was recrystallized from chloroform. ¹H NMR (500 MHz, CDCl₃) δ 8.91 (s, 8H); 8.22 (d, 4H, J=6.5); 7.75 (m, 6H); 7.59 (d, 2H, J=2.2); 7.26 (broad s, 4H); 3.86 (s, 6H); 3.54 (s, 6H). ¹³C NMR (125 MHz, CDCl₃) δ 154.10(0), 152.30(0), 150.13(0), 143.00(0), 134.10(1), 132.62(0), 132.00(1), 131.44(1), 127.35(1), 126.44(1), 121.34(1), 120.69(0), 116.59(0), 114.76(1), 112.31(1), 56.70(3), 55.95(3). Vis - 424 (5.64), 551 (4.34), 584 (3.43).

The C_{2v} isomer followed the C_{2h} isomer off the column. The solvent was evaporated leaving 30 mg (32%) of pure 5,15-bis(2,5-dimethoxyphenyl)-10,20-diphenylporphyrinato zinc. This compound is much more soluble in chloroform than the C_{2h} isomer. The assignment of stereochemistry was made from the NMR data. ¹H NMR (500 MHz, CDCl₃) δ 8.90 (s, 8H); 8.21 (d, 2H, J=7.9); 8.19 (d, 2H, J=6.5); 7.73 (m, 6H); 7.58 (s, 2H); 7.24 (broad s, 4H); 3.84 (s, 6H); 3.53 (s, 6H). ¹³C NMR (125 MHz, CDCl₃) δ 154.14(0); 152.31(0), 150.15(0), 142.94(0), 134.40(1), 132.66(0), 132.02(1), 131.48(1), 127.37(1), 126.46(1), 126.44(1), 121.30(1), 120.72(0), 116.69(0), 114.73(1), 112.28(1), 56.75(3), 55.92(3).

D. 5,15-Bis[(4-methyl)-4'-methyl-2,2'-dipyridyl]-10,20-diphenylporphyrinato zinc

The organometallic reagent was tri-*n*-butyl[(4-methyl)-4'-methyl-2,2'-dipyridyl] tin, prepared by lithiating

-21-

4,4'-dimethyl-2,2'-dipyridyl with one equivalent of lithium diisopropylamide in THF at -78°C and warming the reaction mixture to room temperature. The organolithium reagent was treated with 1.1 equivalent of tributyltin chloride. The
5 resulting organotin reagent was used without further purification.

Chromatography of the crude reaction mixture was carried out on silica with a mixture of CH₂Cl₂, isopropanol, and triethylamine. The porphyrin was eluted in one broad
10 band. The product obtained from this procedure (68% yield) was contaminated with a small amount (0.2 eq per eq of porphyrin) of triphenylphosphine. ¹H NMR (500 MHz, CDCl₃) δ
9.37 (d, 4H, J=4.7); 8.87 (d, 4H, J=4.7); 8.52 (s, 2H); 8.29 (d, 2H, J=5.1); 8.20 (d, 2H, J=5.2); 8.10 (m, 6H); 7.71 (m,
15 6H); 7.01 (d, 2H, J=5.0); 6.88 (d, 2H, J=4.2); 6.46 (s, 4H); 2.32 (s, 6H).

E. 5,15-Bis(trimethylsilylethynyl)-10,20-diphenylporphyrinato zinc

The organometallic reagent was trimethylsilyl
20 ethynyl zinc chloride prepared from trimethylsilylethynyl lithium and anhydrous zinc chloride in THF.

After 48 hours the reaction was bright green. The crude solid was absorbed on silica gel, loaded onto a column packed in hexane, and chromatographed with 20%-30%
25 CH₂Cl₂/hexane. Clean separation of the product from the small quantities of deprotected products were obtained by this method. The solvents were evaporated and the purple solid was washed twice with hexane and dried in vacuo. ¹H NMR (500 MHz, CDCl₃) δ 9.68 (d, 4H, J=4.6); 8.89 (d, 4H, J=4.6); 8.15 (dd,
30 4H, J₁=7.9, J₂=1.7); 7.75 (m, 6H); 0.58 (s, 18H). ¹³C NMR (125 MHz, CDCl₃) δ 152.22, 150.26, 142.10, 134.39, 132.77, 131.29, 127.69, 126.67, 115.08, 107.34, 102.00, 0.32.

EXAMPLE 5

PYRROLE-SUBSTITUTED PORPHYRINS

35 General Procedure

2-Bromo-5,10,15,20-tetraphenylporphyrinato zinc (0.1 mmol) and palladium 1,1'-bis(diphenylphosphino)ferrocene)

-22-

dichloride (Pd(dppf)Cl_2 , 7 mg) were combined with 1.0 mmol of the organometallic reagent indicated below in 35 ml dry, degassed THF. The solution was allowed to stand for 12 hours, the solvent evaporated, and the compound purified by flash chromatography.

A. 2-Vinyl-5,10,15,20-tetraphenyl porphyrinato zinc

The organometallic reagent was tributylvinyl tin.

The crude reaction mixture was chromatographed on silica and eluted with 50% CH_2Cl_2 /hexane. ^1H NMR (250 MHz, CDCl_3) δ 8.97 (s, 1H); 8.90 (m, 4H); 8.87 (d, 1H, $J=4.7$); 8.79 (d, 1H, $J=4.7$); 8.20 (m, 6H); 8.06 (d, 2H, $J=6.6$); 7.74 (m, 12H); 6.39 (dd, 1H, $J_1=17.0$, $J_2=9.1$); 5.83 (dd, 1H, $J_1=17.1$, $J_2=2.0$); 5.01 (dd, 1H, $J_1=10.7$, $J_2=2.0$). Vis (CHCl_3) 426 (5.53), 517 (3.68); 555 (4.22), 595 (3.68).

B. 2-(2,5-dimethoxyphenyl)-5,10,15,20-tetraphenyl porphyrinato zinc

The organometallic reagent was 2,5-dimethoxyphenyl zinc chloride, prepared from the corresponding lithium reagent and anhydrous zinc chloride in THF/diethyl ether.

Flash chromatograph of the crude reaction mixture was carried out with chloroform. The title compound was isolated in 78% yield. ^1H NMR (500 MHz, CDCl_3) δ = 8.94 (d, 1H, $J=4.7$); 8.93 (s, 2H); 8.92 (d, 1H, $J=4.8$); 8.85 (s, 1H); 8.84 (d, 1H, $J=4.7$); 8.70 (d, 1H, $J=4.7$); 8.23 (m, 6H); 7.98 (d, 1H, $J=7.0$); 7.70 (m, 10H); 7.25 (quintet, 2H, $J=7.4$); 7.15 (t, 1H, $J=7.0$); 6.92 (d, 2H, $J=3.1$); 6.54 (dd, 1H, $J_1=9.0$, $J_2=3.2$); 6.40 (d, 1H, $J=9.1$); 3.68 (s, 3H); 3.42 (s, 3H). ^{13}C NMR (125 MHz, CDCl_3) δ = 152.59(0), 151.33(0), 150.46(0), 150.31(0), 150.27(0), 150.15(0), 150.12(0), 150.03(0), 148.30(0), 147.16(0), 143.32(0), 142.97(0), 142.86, 140.71(0), 135.63(1), 135.20(1), 134.45(1), 134.13(1), 132.52(1), 132.02(1), 131.91(1), 131.82(1), 131.32(1), 129.27(0), 127.44(1), 127.38(1), 127.18(1), 126.53(1), 126.50(1), 124.91(1), 124.70(1), 122.36(0), 121.30(0), 120.91(0), 120.54(0), 118.15(1), 113.03(1), 110.35(1), 55.98(3).

-23-

54.87(3). Vis (CHCl₃) 421.40 (5.60), 513.2 (3.45), 549.75 (4.28), 587.15 (3.45).

C. 2-(Trimethylsilyl ethynyl)-5,10,15,20-tetraphenyl porphyrinato zinc

5 The organometallic reagent, trimethylsilylacetylide zinc chloride, was prepared from the corresponding lithium reagent and anhydrous zinc chloride in THF.

The crude reaction mixture was chromatographed on silica and eluted with 50% CH₂Cl₂/hexane. ¹H NMR (250 MHz, CDCl₃) δ 9.25 (s, 1H); 8.89 (m, 4H); 8.85 (d, 1H, J=4.9); 8.76 (d, 1H, J=4.9); 8.16 (m, 6H); 8.09 (d, 2H, J=7.1); 7.67 (m, 12H); 0.21 (s, 9H). Vis (CHCl₃) 431 (5.43), 523 (shoulder); 559 (4.18), 598 (3.67).

15 D. 2-n-butyl-5,10,15,20-tetraphenyl porphyrinato zinc

Butylzinc chloride was prepared from n-butyllithium and anhydrous zinc chloride in THF.

The crude reaction mixture was chromatographed on silica and eluted with 50% CH₂Cl₂/hexane. ¹H NMR (250 MHz, CDCl₃) δ 8.97 (m, 4H); 8.91 (d, 1H, J=4.7); 8.77 (d, 1H, J=4.7); 8.74 (s, 1H); 8.22 (m, 6H); 8.13 (d, 2H, J=7.3); 7.77 (m, 12H); 2.81 (t, 2H, J=7.7); 1.83 (quint, 2H, J=7.8); 1.30 (quint, 2H, J=7.6); 0.91 (t, 3H, J=8.2).

EXAMPLE 6

25 VINYLIC-BRIDGED PORPHYRINS AND THEIR POLYMERS

A. cis-Bis-1,2-[5-(10,15,20-triphenylporphyrinato) zinc] ethene

5-Bromo-10,15,20-triphenylporphyrinato zinc (0.2 mmol) and Pd(PPh₃)₄ (0.02 mmole) are dissolved in 25 ml dry, degassed THF. A solution of cis-bis(tri-n-butyltin)ethene (0.2 mmol) in 5 ml THF is added and the solution heated at reflux for 2 days. The reaction is quenched with water, extracted with methylene chloride, dried over calcium chloride, and the solvents are evaporated. The crude solid is chromatographed on silica using methylene chloride/hexane

-24-

eluant to isolate a dimer having formula (3), wherein R_{A1} , R_{A2} , and R_{A4} are phenyl and M is Zn.

5 B. *cis*-Bis-1,2-{5-[10,15,20-tris(pentafluorophenyl)]-2,3,7,8,12,13,17,18-octakis-(trifluoromethyl)porphyrinato zinc}-1,2-difluoroethene


5-Bromo-10,15,20-tris(pentafluorophenyl)porphyrinato zinc (0.2 mmol) and $Pd(PPh_3)_4$ (0.02 mmol) are dissolved in 25 ml dry THF. A solution of *cis*-bis(tri-*n*-butyltin)-1,2-difluoroethene (0.2 mmol) in 5 ml THF is added and the solution heated at reflux for 2 days. The reaction is quenched with water, extracted with methylene chloride, dried over calcium chloride, and the solvents evaporated. The crude solid is chromatographed on silica using methylene chloride/hexanes eluent to isolate *cis*-bis-1,2-{5-[10,15,20-tris(pentafluorophenyl)porphyrinatozinc]-1,2-difluoroethene. This material is dissolved in chloroform and reacted with a large excess of N-bromosuccinimide as in Example 2 to perbrominate positions R_{B1} - R_{B8} on both porphyrins. The resulting material filtered through a fine fritted disk and washed with water, methanol, and acetone, dried in vacuo, and then recrystallized from THF/heptane. *cis*-Bis-1,2-{5-[10,15,20-tris(pentafluorophenyl)-2,3,7,8,12,13,17,18-octabromoporphyrinato zinc is reacted with $Pd(dppf)$ and a large excess of $CuCF_3$ in the dark as in Example 4. After a reaction time of about 48 hours, the product is chromatographed on silica with CH_2Cl_2/CCl_4 eluent to yield the title compound.

30 C. Cofacial-bis-[*cis*-ethenyl meso-bridged] porphyrin [CEBP] (Formula (5)) and Polymeric-bis-[*cis*-ethenyl meso-bridged] porphyrin [PABP] (Formula (6))


5,15-Dibromo-10,20-diphenylporphyrinato zinc (0.2 mmol) and $Pd(PPh_3)_4$ (0.02 mmole) are dissolved in 25 ml dry, degassed THF. A solution of *cis*-bis(tri-*n*-butyltin)ethene (0.2 mmol) in 5 ml THF is added and the solution heated at reflux for 2 days. The reaction is quenched with water,

-25-

extracted with methylene chloride, dried over calcium chloride, and the solvents are evaporated. The crude solid is chromatographed on silica using methylene chloride/hexane eluant to isolate the Cofacial-bis-[*cis*-ethenyl *meso*-bridged] zinc porphyrin complex of formula (5) and Polymeric-bis-[*cis*-ethenyl *meso*-bridged] porphyrin species of formula (6), wherein R_{A1} and R_{A3} are phenyl and M is Zn.



(5)



(6)

-26-

D. Fluorinated Cofacial-bis-[*cis*-ethenyl meso-bridged] porphyrin [FCEBP] and Fluorinated Polymeric-bis-[*cis*-ethenyl meso-bridged] porphyrin [FPEBP]


5 5,15-Dibromo-10,20-bis(pentafluorophenyl) porphyrinato zinc (0.2 mmol) and Pd(PPh₃)₄ (0.02 mmol) are dissolved in 25 ml dry THF. A solution of *cis*-bis(tri-*n*-butyltin)-1,2-difluoroethene (0.02 mmol) in 5 ml THF is added and the solution heated at reflux for 2 days. The reaction
10 is quenched with water, extracted with methylene chloride, dried over calcium chloride, and the solvents evaporated. The crude solid is chromatographed on silica using methylene chloride/hexanes eluent to isolate the Cofacial-bis-[*cis*-ethenyl meso-bridged] zinc porphyrin complex as well as the
15 Polymeric-bis-[*cis*-ethenyl meso-bridged] porphyrin species. The cofacial and polymeric species are dissolved separately in chloroform. The cofacial porphyrin complex dissolved in chloroform and reacted with a large excess of *N*-bromosuccinimide as in Example 2 to perbrominate positions
20 R_{B1}-R_{B8} on both porphyrins. The resulting material filtered through a fine fritted disk and washed with water, methanol, and acetone, dried in vacuo, and then recrystallized from THF/heptane to yield the title compound. The isolated material is reacted with Pd(dppf) and a large excess of CuCF₃
25 in the dark in a manner as in Example 4. After a reaction time of about 48 hours, the product is chromatographed on silica with CH₂Cl₂/CCl₄ eluent to yield a perfluorinated CEPB analogous to formula (5). Perfluorinated PEBP is synthesized in a similar manner, yielding a species analogous to formula
30 (6) where highly fluorinated porphyrins are linked via fluorovinyl units.

E. Cofacial-bis-[1,8-anthracenyl-meso-bridged] porphyrin [CBAP] (Formula (7)) and Polymeric-bis-[1,8-anthracenyl-meso-bridged] [PBAP]
35 porphyrin (Formula (8))

5,15-(Dibromoporphyrinato)zinc (0.2 mmol) and Pd(PPh₃)₄ (0.02 mmol) are dissolved in 25 ml dry, degassed


-27-

THF. A solution of 1,8-anthracenyl-bis-(tributyl tin) (0.2 mmol) in 5 ml THF is added and the solution heated at reflux for 2 days. The reaction is quenched with water, extracted with methylene chloride, dried over calcium chloride, and the
5 solvents are evaporated. The crude solid is chromatographed on silica using methylene chloride/hexane eluant to isolate the Cofacial-bis-[1,8-anthracenyl-meso-bridged] zinc porphyrin complex of formula (7) and the Polymeric-bis-[1,8-anthracenyl-meso-bridged] zinc porphyrin species of formula (8), where R_{A1}
10 and R_{A3} are phenyl and M is Zn.



(7)

-28-

**EXAMPLE 7****ACETYLENIC PORPHYRIN POLYMERS**

- 5 A. Poly(5,15-bis(ethynyl)-10,20-diphenylporphyrinato zinc)

5,15-Bis(ethynyl)-10,20-diphenylporphyrinato zinc (0.2 mmol) in pyridine (20 ml) is slowly added to a solution of cupric acetate (0.4 mmol) in 20 ml 1:1 pyridine/methanol generally according to the procedure of Eglinton, et al., The
 10 Coupling of Acetylenic Compounds, p. 311 in Advances in Organic Chemistry, Raphael, et al., eds., 1963, Interscience Publishers.

- B. Poly(5,15-bis(ethynylphenyl)-10,20-diphenylporphyrinato zinc)

-29-

5,15-Diethynyl-10,20-diphenylporphinato zinc (0.2 mmol) and 1,4-dibromobenzene are combined in a mixture of 30 ml toluene and 10 ml diisopropylamine. CuI (0.4 mmol) and Pd(Ph₃)₄ (0.02 mmol) are added and the mixture is heated at 65°C for 3 days. The crude solid is washed with methanol and acetone and dried in vacuo.

Alternatively, the polymer is prepared from 1,4-diethynylbenzene and 5,15-dibromo-10,20-diphenylporphinato zinc via the identical procedure.

10 EXAMPLE 8

DOPED PORPHYRIN POLYMERS

5,15-Bis(ethynyl)-10,20-diphenylporphyrinato zinc is polymerized according to the generally procedure provided by Skotheim, ed., Handbook of Conducting Polymers, Volume 1, pp. 405-437, Marcel Dekker, 1986 using a catalytic amount of MoCl₅, Mo(CO)₆, WCl₆, or W(CO)₆. The resultant polymer is then doped with an oxidant such as iodine or SbF₅.

EXAMPLE 9

METALATION OF BROMINATED PORPHYRINS

Finely divided zinc metal was prepared generally according to the method of Rieke (*J. Org. Chem.* 1984, 49, 5280 and *J. Org. Chem.* 1988, 53, 4482) from sodium naphthalide and zinc chloride (0.18 mmol each) in THF. A solution of [5-bromo-10,20-diphenylporphinato]zinc (100 mg, 0.17 mmol) dissolved in 40 mL THF was added by syringe to the zinc metal suspension, and the mixture was stirred at room temperature overnight; during this time all of the zinc metal dissolved. The ring-metalated porphyrin is suitable for palladium-catalyzed coupling with a variety of aryl and vinyl halides.

30 EXAMPLE 10

PALLADIUM-CATALYZED CROSS-COUPLING WITH RING-METALATED PORPHYRINS

5-[(10,20-Diphenylporphinato)zinc]zincbromide (0.2 mmol) is prepared in 15 mL of THF as in Example 9, above and

-30-

is placed in a dry 100 mL Schlenk tube. A solution of 2-iodothiophene (0.4 mmol) in 5 mL of THF is added via syringe. Pd(dppf) (3 mg) is prepared by stirring a suspension of Pd(dppf)Cl₂ in THF over Mg turnings for 20 min. and is transferred into the reaction mixture by canula. The solution is stirred overnight, quenched with aqueous ammonium chloride, extracted with CH₂Cl₂, and dried over CaCl₂. The solvent is evaporated to dryness and chromatography is carried out with 1:1 CH₂Cl₂ as eluant. The product, [5-(2-thiophenyl)-10,20-diphenylporphinato]zinc, elutes in one band and is isolated in 90% yield.

EXAMPLE 11**POLYMERIZATION WITH RING-METALATED PORPHYRIN DERIVATIVES**

[5,15-Bis(zinc bromide)-10,20-diphenylporphinato]-zinc (0.2 mmol) is prepared in 15 mL of THF as in Example 9 and is placed in a dry 100 mL Schlenk tube. A solution of [5,15-dibromo-10,20-diphenylporphinato]zinc (0.2 mmol) in 15 mL of THF is added by syringe. Pd(dppf) (3 mg) is prepared by stirring a suspension of Pd(dppf)Cl₂ in THF over Mg turnings for 20 min. and is transferred into the reaction mixture by canula. The mixture is heated at 60°C for 3 days, cooled to room temperature and filtered through a fine-fritted glass disk. The filtered polymer is washed with hexane followed by methanol and dried in vacuo.

EXAMPLE 12**CARBONYLATION OF [5-BROMO-10,20-DIPHENYLPORPHINATO]ZINC**

5-[(10,20-Diphenylporphinato)zinc]magnesium bromide (0.2 mmol) is prepared in 15 mL of THF as in Example 9 and is placed in a dry 100 mL Schlenk tube. The vessel is cooled to 0°C and dry CO₂ gas is bubbled through the solution. The solution is stirred for 1 h at room temperature, quenched with 0.1 M HCl, extracted with CH₂Cl₂, and dried over CaCl₂. The solvent is evaporated to dryness and chromatography is carried out with THF:CH₂Cl₂ as eluant. Upon evaporation of the

-31-

solvent [5-carboxy-10,20-diphenylporphinato]zinc is isolated in 85% yield.

EXAMPLE 13**COUPLING ON UNMETALATED PORPHYRIN DERIVATIVES****5 A. Using Organozinc Chloride Reagents**

Trimethylsilylacetylene (3 mmol) was deprotonated with n-butyl lithium (3 mmol) at -78°C in THF and warmed slowly to room temperature. Excess ZnCl₂ (650 mg) in 5 mL of THF was transferred into the solution via canula. Pd(dppf) (3
10 mg) was prepared by stirring a suspension of Pd(dppf)Cl₂ in THF over Mg turnings for 20 min. and transferred into the solution by canula. The entire reaction mixture was transferred to a dry 100 mL Schlenk tube containing 340 mg of 5,15-dibromo-10,20-diphenylporphyrin. The solution was heated
15 to 40°C and left sealed overnight. TLC of the reaction mixture after 18 h shows a mixture of fluorescent products. The mixture was quenched with aqueous ammonium chloride, extracted with CH₂Cl₂, and dried over CaCl₂. The solvent was evaporated to dryness and chromatography was carried out with
20 1:1 CH₂Cl₂:hexane as eluant. The majority of the material was collected in two bands which proved to be [5-(2-trimethylsilylethynyl)-10,20-diphenylporphinato]zinc and [5,15-bis(2-trimethylsilylethynyl)-10,20-diphenylporphinato]-zinc. The two products were isolated in 83% overall yield.

25 B. Using Organotrialkyltin Reagents

5,15-Dibromo-10,20-diphenylporphyrin is placed in a dry 100 mL Schlenk tube and dissolved in 30 mL of THF. A solution of vinyltributyltin (3 mmol) in 5 mL THF is added to the reaction mixture. Pd(dppf) (3 mg) is prepared by stirring
30 a suspension of Pd(dppf)Cl₂ in THF over Mg turnings for 20 min. and is transferred into the reaction mixture by canula. The solution is stirred overnight, quenched with aqueous ammonium chloride, extracted with CH₂Cl₂, and dried over CaCl₂. The solvent is evaporated to dryness and chromatography is
35 carried out with 1:1 CH₂Cl₂:hexane as eluant. The product,

-32-

5,15-diphenyl-10,20-divinylporphyrin, elutes in one band and is isolated in 90% yield.

EXAMPLE 14**COUPLING ON DILITHIATED PORPHYRIN DERIVATIVES.**

5 A solution of N,N"-dilithio-5,15-dibromo-10,20-diphenylporphyrin (0.2 mmol) in 15 mL of THF is prepared generally according to the method of Arnold, *J. Chem. Soc. Commun.* 1990, 976. A solution of vinyltributyltin (2 mmol) in 5 mL THF is added to the reaction mixture. Pd(dppf) (3 mg)
10 is prepared by stirring a suspension of Pd(dppf)Cl₂ in THF over Mg turnings for 20 min. and is transferred into the reaction mixture by canula. The solution is stirred overnight, and quenched with a solution of anhydrous NiCl₂ in THF. Aqueous ammonium chloride is added, the solution is
15 extracted with CH₂Cl₂, and dried over CaCl₂. The solvent is evaporated to dryness and chromatography is carried out with 1:1 CH₂Cl₂:hexane as eluant. The product, [5,15-diphenyl-10,20-divinylporphinato]nickel, elutes in one band and is isolated in 90% yield.


20 Those skilled in the art will appreciate that numerous changes and modifications may be made to the preferred embodiments of the invention and that such changes and modifications may be made without departing from the spirit of the invention. For example, it is believed that the
25 methods of the present invention can be practiced using porphyrin-related compounds such as chlorins, phorbins, bacteriochlorins, porphyrinogens, sapphyrins, texaphrins, and phthalocyanines in place of porphyrins. It is therefore intended that the appended claims cover all such equivalent
30 variations as fall within the true spirit and scope of the invention.

-33-


WHAT IS CLAIMED IS:

1. A process for preparing a substituted porphyrin, comprising the steps of:


providing a porphyrin compound having formula (1), (2), or (3):



(1)



(2)



(3)

wherein M and M' are metal atoms and at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} is a first halogen;

preparing a ring-metalated first reaction product by a process selected from:

contacting said porphyrin compound with an activated metal Z under conditions effective to ring-metalate said porphyrin compound; or

contacting said porphyrin compound with an alkyltin reagent under conditions effective to ring-metalate said porphyrin compound; or

-34-

contacting said first reaction product with a compound having formula R_p-R_z , where R_p is cyclopentadienyl or aryl having about 6 to about 20 carbon atoms, alkyl having 1 to about 10 carbon atoms, alkenyl having 2 to about 10 carbon atoms, or porphyrinato and R_z is a leaving group, said contacting being performed in the presence of a complex having formula $Y(L)$, wherein Y is a second metal and L is a ligand, under conditions effective to produce a substituted porphyrin.

2. The process of claim 1 wherein said halogen is Cl, Br, or I.

3. The process of claim 1 wherein Y is Pd, Ni, Pt, Ru, or Cu.

4. The process of claim 1 wherein Y is Pd or Ni and L is a alkylphosphine, arylphosphine, alkylarsenine, a arylarsenine, alkylstibene, arylstibene, or nitrogen-containing heterocycle.

5. The process of claim 1 wherein said first reaction product includes a compound having formula P_N-Z-X_A wherein P_N is said porphyrin compound and X_A is said halogen.

6. The process of claim 1 wherein Z is Zn, Mg, Ca, Cd, Cu, Sn, Sr, Ba, or Zr.

7. The process of claim 1 wherein R_p is aryl having about 6 to about 20 carbon atoms, alkyl having 1 to about 10 carbon atoms, or alkenyl having 2 to about 10 carbon atoms.

8. The process of claim 1 wherein R_z is halogen, tosylate, or triflate.

9. The process of claim 1 wherein said activated metal is produced by contacting a metal reagent having formula

-35-

$Z(X_c)_c$, wherein X_c is a halogen and c is 1-4, with a reducing agent.

10. The process of claim 1 wherein said porphyrin compound and said alkyltin reagent are contacted in the presence of a complex having formula $Y(L)_2$.


11. The process of claim 1 wherein said alkyltin reagent has formula $(R)_3Sn-Sn(R)_3$, where R is alkyl having 1 to about 10 carbon atoms.

~~12. The product of the process of claim 1.~~


13. A process for derivatizing a porphyrin, comprising the steps of:

providing a porphyrin compound having formula (1), (2), or (3):


-36-



(1)



(2)



(3)

wherein M and M' are metal atoms and at least one of R_{A1} - R_{A4} or R_{B1} - R_{B4} is a halogen; and

contacting said porphyrin compound with a derivatizing agent wherein:

said derivatizing agent is an activated metal Z; or
said derivatizing agent is an alkyltin reagent.

14. The process of claim 13 wherein said activated metal is produced by contacting a metal reagent having formula $Z(X_c)_c$, wherein X_c is a halogen and c is 1-4, with a reducing agent.

15. The process of claim 13 wherein said porphyrin compound and said activated metal are contacted under conditions that produce a first derivatized porphyrin having

-37-

formula P_N-Z-X_A wherein P_N is said porphyrin compound and X_A is said halogen.

16. The process of claim 13 wherein said porphyrin compound and said alkyltin reagent are contacted in the presence of a complex having formula $Y(L)_2$.

17. The process of claim 13 wherein said alkyltin reagent has formula $(R)_3Sn-Sn(R)_3$, where R is alkyl having 1 to about 10 carbon atoms.

18. ~~The process of claim 17 wherein said porphyrin compound and said alkyltin reagent are contacted under conditions that produce a first derivatized porphyrin having formula $P_N-Sn(R)_3$, wherein P_N is said porphyrin compound.~~

19. The product of the process of claim 13.


20. The process of claim 13 further comprising contacting said first derivatized porphyrin with a quenching reagent.

21. The process of claim 20 wherein said quenching reagent is selected from CO_2 , esters, aldehydes, ketones, acid chlorides, and amides.


22. The process of claim 20 wherein said quenching reagent is selected from trialkyl tin halides and trialkyl borates.

-38-


23. A compound having formula (1), (2), or (3):



(1)



(2)




(3)

wherein at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $-Z-X_A$ or $-\text{Sn}(R)$, where M and M' are metal atoms, Z is a metal, X_A is a halogen, and R is alkyl having 1 to about 10 carbon atoms.


24. The compound of claim 23 wherein Z is Zn, Mg, Ca, Cd, Cu, Sn, Sr, Ba, or Zr.

-39-


25. A polymer comprising a plurality of linked porphyrin units having formula (1), (2), or (3):



(1)



(2)




(3)


herein M and M' are metal atoms and at least two of said porphyrin units share a covalent bond.

-40-

26. A compound having formula (1) or (2):



(1)



(2)

wherein:

- (a) at least one of R_{A1} - R_{A4} has formula $CH=CH_2$; or
- (b) at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C(R_C)=C(R_D)(R_E)$, provided that at least one of R_C , R_D , and R_E is not H; or
- (c) at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C\equiv C(R_D)$; or
- (d) at least one of R_{A1} - R_{A4} is haloalkyl having 1 to about 20 carbon atoms; or
- (e) at least one of R_{B1} - R_{B8} is haloalkyl having 2 to about 20 carbon atoms; or
- (f) at least five of R_{B1} - R_{B8} are haloalkyl having 1 to about 20 carbon atoms or haloaryl having about 6 to about 20 carbon atoms; or
- (g) at least one of R_{B1} - R_{B8} is haloaryl or haloheteroaryl having about 6 to about 20 carbon atoms;

where R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 6 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato, or a chemical functional group comprising a peptide, nucleoside or saccharide; and

M is a chelated metal atom.

-41-

27. The compound of claim 26 wherein at least one of R_{A1} - R_{A4} has formula $CH=CH_2$.

28. The compound of claim 26 wherein at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C(R_C)=C(R_D)(R_E)$, provided that at least one of R_C , R_D , and R_E is not H.

29. The compound of claim 26 wherein at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C\equiv C(R_D)$.

30. The compound of claim 26 wherein at least one of R_{A1} - R_{A4} is haloalkyl having 1 to about 20 carbon atoms.

31. The compound of claim 26 wherein at least one of R_{B1} - R_{B8} is haloalkyl having 2 to about 20 carbon atoms.

32. The compound of claim 26 wherein at least five of R_{B1} - R_{B8} are haloalkyl having from 1 to about 20 carbon atoms.

33. The compound of claim 26 wherein at least one of R_{B1} - R_{B8} is haloaryl having about 6 to about 20 carbon atoms.

34. The compound of claim 26 wherein at least one of R_{B1} - R_{B8} is haloaryl or haloheteroaryl having about 6 to about 20 carbon atoms.

35. The compound of claim 26 wherein at least one of R_{B1} - R_{B8} is alkyl having 1 to about 20 carbon atoms or aryl having 6 to about 20 carbon atoms.

36. The compound of claim 36 wherein R_{B5} has formula $C\equiv C(R_D)$ and each of R_{A1} - R_{A4} are alkyl having 1 to about 20 carbon atoms or aryl having 6 to about 20 carbon atoms.

37. The compound of claim 36 wherein R_{B1} and R_{B5} have formula $C\equiv C(R_D)$ and each of R_{A1} - R_{A4} are alkyl having 1 to

-42-

about 20 carbon atoms or aryl having 6 to about 20 carbon atoms.

38. The compound of claim 36 wherein R_{B2} and R_{B5} have formula $C\equiv C(R_D)$ and each of R_{A1} - R_{A4} are alkyl having 1 to about 20 carbon atoms or aryl having 6 to about 20 carbon atoms.

39. The compound of claim 36 wherein each of R_{B1} - R_{B8} have formula $C\equiv C(R_D)$ and each of R_{A1} - R_{A4} are alkyl having 1 to about 20 carbon atoms or aryl having 6 to about 20 carbon atoms.

40. The compound of claim 36 wherein at least one of R_{B1} - R_{B8} is H, alkyl having 1 to about 20 carbon atoms, aryl having 6 to about 20 carbon atoms, F, Cl, Br, or I.

41. The compound of claim 40 wherein R_{A1} has formula $C\equiv C(R_D)$, each of R_{A2} - R_{A4} are alkyl having 1 to about 20 carbon atoms, aryl having 6 to about 20 carbon atoms, F, Cl, Br, or I, and R_D is H, F, Cl, Br, I, alkyl having from 1 to about 20 carbon atoms, aryl having about 6 to about 20 carbon atoms, alkenyl having from 1 to about 20 carbon atoms, alkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide.

42. The compound of claim 40 wherein R_{A1} and R_{A3} have formula $C\equiv C(R_D)$, R_{A2} and R_{A4} are alkyl having 1 to about 20 carbon atoms, aryl having 6 to about 20 carbon atoms, F, Cl, Br, or I, and R_D is H, F, Cl, Br, I, alkyl having from 1 to about 20 carbon atoms, aryl having about 6 to about 20 carbon atoms, alkenyl having from 1 to about 20 carbon atoms, alkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide.

-43-

43. The compound of claim 42 wherein R_p is 3,4,5-trialkyl-substituted phenyl or 3,4,5-trialkyloxy-substituted phenyl wherein said alkyl and alkoxy substituents have about 7 to about 20 carbon atoms.

44. The compound of claim 40 wherein $R_{A1}-R_{Am}$ have formula $C\equiv C(R_p)$ and R_p is H, F, Cl, Br, I, alkyl having from 1 to about 20 carbon atoms, aryl having about 6 to about 20 carbon atoms, alkenyl having from 1 to about 20 carbon atoms, alkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide.

45. The compound of claim 44 wherein R_p is 3,4,5-trialkyl-substituted phenyl or 3,4,5-trialkyloxy-substituted phenyl wherein said alkyl and alkoxy substituents have about 7 to about 20 carbon atoms.

46. The compound of claim 26 wherein M is a lanthanide or actinide or Sc, Ti, V, Cr, Mn, Fe, Co, Ni, Cu, Y, Zr, Nb, Mo, Ru, Rh, Pd, Ag, La, Hf, Ta, W, Re, Os, Ir, Pt, Cd, Hg, Li or Au.


47. The compound of claim 26 wherein M is a metal having a full valence shell.

48. The compound of claim 26 wherein M is Zn or Mg.


-44-

49. A polymer comprising a plurality of linked porphyrin units, wherein:

each of said porphyrin units, independently, has formula (1) or (2):



(1)



(2)

wherein:

at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} includes a linking group that is:

$[C(R_C)=C(R_D)(R_E)]_x$, $[C\equiv C(R_D)]_x$, $[CH_2(R_C)-CH(R_D)(R_E)]_x$ or $[CH=CH(R_D)]_x$ where n is at least 1, where R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 6 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide; or

cycloalkyl or aryl having about 10 to about 22 carbon atoms; and


M is a chelated metal atom;

50. The polymer of claim 49 wherein the remaining of R_{A1} - R_{A4} and R_{B1} - R_{B8} are, independently, H, alkyl or heteroalkyl having 1 to about 20 carbon atoms, aryl or heteroaryl having 6 to about 20 carbon atoms, $C(R_C)=C(R_D)(R_E)$,

-45-


$C\equiv C(R_p)$, or a chemical functional group comprising a peptide, nucleoside or saccharide.

51. The polymer of claim 49 wherein a portion of said polymer has formula:



wherein Q_L is a linking group, P_N is a porphyrin unit, and h , 1, and s are, independently, at least 1.

52. The polymer of claim 49 wherein a portion of said polymer has formula:




wherein Q_H is a linking group, and h , u , and v are, independently, at least 1.

53. The polymer of claim 49 wherein at least two of said porphyrin units are covalently bound with a polyacrylate, polyolefin, polyether, polyurethane, polycarbonate, polyaniline, polypyrrole, polythiophene, or polyacetylene.

54. A process for preparing a substituted porphyrin, comprising the steps of:

providing a porphyrin compound having formula (2):



(2)

wherein M is a chelated metal atom and at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} is a first halogen;

contacting said porphyrin compound with a

-46-

complex having formula $Y(L)_2$, wherein Y is a first metal and L is a ligand, said contacting being performed under conditions effective to produce a first reaction product;

contacting said first reaction product with an organometallic compound having formula $T(R_L)_z(R_O)_t$, $T(R_L)_z(R_O)_t(X_B)_w$, $T(R_O)(X_B)$ or $T(R_O)_t$ where:

T is a second metal;

X_B is a second halogen or an alkoxy group;

R_L is cyclopentadienyl or aryl having about 6 to about 20 carbon atoms;

R_O is alkyl, alkenyl or alkynyl having 1 to about 10 carbon atoms, aryl having about 6 to about 20 carbon atoms; and

z and w are greater than or equal to 0 and t is at least 1;

said contacting being performed under conditions effective to produce a second reaction product comprising a substituted porphyrin having formula (1) or (2) wherein at least one of at least one of $R_{A1}-R_{A4}$ or $R_{B1}-R_{B8}$ is R_O .

55. The process of claim 49 wherein said first halogen is Cl or Br.

56. The process of claim 54 wherein Y is Pd, Ni, Pt, Ru, or Cu.

57. The process of claim 54 wherein Y is Pd or Ni and L is an alkylphosphine, arylphosphine, or a nitrogen-containing heterocycle.

58. The process of claim 54 wherein said first reaction product includes a compound having formula $P_N-Y(L)_2X_A$ wherein P_N is said porphyrin compound and X_A is said first halogen.


59. The process of claim 54 wherein T is Li, Na, K, Rb, Cs, Hg, Sn, Al, B, Si, Zn, Zr, Cd, Cu, or Mg.

-47-

60. The process of claim 54 wherein said second reaction product includes a compound having formula $P_N-Y(L)_2R_0$ wherein P_N is said porphyrin compound.

61. The process of claim 54 wherein R_0 is $C(R_C)=C(R_D)(R_E)$, $C\equiv C(R_D)$, haloalkyl having from 1 to about 20 carbon atoms, or haloaryl or haloheteroaryl having about 6 to about 20 carbon atoms, and R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 6 to about 20 carbon atoms, ~~alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide.~~

62. A process for derivatizing a porphyrin compound, comprising the steps of:
providing a porphyrin compound having formula (2):



(2)

wherein at least one of $R_{A1}-R_{A4}$ or $R_{B1}-R_{B8}$ is a first halogen;
and M is chelated metal atom; and

contacting said porphyrin compound with a complex having formula $Y(L)_2$, where Y is a metal and L is a ligand, said contacting being performed under conditions effective to produce a first reaction product that includes

-48-

a compound having formula $P_N-Y(L)_2X_A$ wherein P_N is said porphyrin compound and X_A is said first halogen.

63. The product of the process of claim 62.

64. The process of claim 62 further comprising contacting said first reaction product with an organometallic compound having formula $T(R_L)_z(R_O)_t(X_B)_w$, $T(R_O)(X_B)$ or $T(R_O)_t$ where:

T is a Li, Zn, Cd, Cu, Sn, Mg or Zr;

X_B is a second halogen or an alkoxy group;

~~R_L is cyclopentadienyl or aryl having about 6~~
to about 20 carbon atoms;

R_O is $C(R_C)=C(R_D)(R_E)$, $C\equiv C(R_D)$, haloalkyl having from 1 to about 20 carbon atoms, or haloaryl or haloheteroaryl having about 6 to about 20 carbon atoms, and R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 6 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide; and


z and w are greater than or equal to 0 and t is at least 1;

said contacting being performed under conditions effective to produce a second reaction product that includes a compound having formula $P_N-Y(L)_2R_O$.


65. The product of the process of claim 62.

-49-

66. A compound having formula (1) or (2):



(1)



(2)

wherein:


at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} is a halogen; and

at least one of the remaining R_{A1} - R_{A4} or R_{B1} - R_{B8} is $C(R_C)=C(R_D)(R_E)$, $C\equiv C(R_D)$, haloalkyl having from 1 to about 20 carbon atoms, or haloaryl or haloheteroaryl having about 6 to about 20 carbon atoms where R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 6 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide; and


M is a chelated metal atom.

-50-

67. A compound having formula (1) or (2):




(1)




(2)

wherein M is a chelated metal atom and at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $Y(L)_2X_A$ where Y is a metal, L is a ligand, and X_A is a halogen.

68. A compound having formula (1) or (2):



(1)



(2)


wherein at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $P_N-Y(L)_2R_0$ where M is a chelated metal atom, Y is a metal; L is a ligand; and R_0 is $C(R_C)=C(R_D)(R_E)$, $C\equiv C(R_D)$, haloalkyl having from 1 to about 20 carbon atoms, or haloaryl or haloheteroaryl having about 6 to about 20 carbon atoms, and R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having

-51-


about 6 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato or a chemical functional group comprising a peptide, nucleoside or saccharide.

69. A process for preparing a polymer, comprising the steps of:

providing at least two compounds that, independently, have formula (1) or (2):



(1)



(2)

wherein:

- (a) at least one of R_{A1} - R_{A4} has formula $CH=CH_2$; or
- (b) at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C(R_C)=C(R_D)(R_E)$, provided that at least one of R_C , R_D , and R_E is not H; or
- (c) at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} has formula $C\equiv C(R_D)$;

where R_C , R_D , and R_E are, independently, H, F, Cl, Br, I, alkyl or heteroalkyl having from 1 to about 20 carbon atoms, aryl or heteroaryl having about 6 to about 20 carbon atoms, alkenyl or heteroalkenyl having from 1 to about 20 carbon atoms, alkynyl or heteroalkynyl having from 1 to about 20 carbon atoms, trialkylsilyl, porphyrinato, or a

-52-

chemical functional group comprising a peptide, nucleoside or saccharide; and

M is a chelated metal atom;

provided that:

- (i) at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} in each of said compounds contains an olefinic carbon-carbon double bond or a chemical functional group reactive therewith; or
 - (ii) at least one of R_{A1} - R_{A4} or R_{B1} - R_{B8} in each of said compounds contains a carbon-carbon triple bond or a chemical functional group reactive therewith; and
-

contacting said compounds for a time and under reaction conditions effective to form covalent bonds through said carbon-carbon double or triple bonds.

70. In a process in which an alkane is selectively oxidized by contact with air or oxygen in the presence of a catalyst comprising a Group IV(a) to VII transition metal coordination complex, the improvement wherein said complex comprises a compound according to claim 26.

71. A supported catalyst comprising a compound according to claim 26 on a solid support material.

72. A method of inactivating infectious agents, comprising the steps of:

forming a mixture by contacting a compound according to claim 1 to a fluid comprising infectious agents; and
subjecting said mixture to irradiation at a wavelength at or near an appropriate low energy absorption maximum of said compound for a time and at an intensity sufficient to inactivate said infectious agents.

73. A method of inactivating infectious agents, comprising the steps of:

-53-

forming a mixture by contacting a polymer according to claim 24 to a fluid comprising infectious agents; and
subjecting said mixture to irradiation at a wavelength at or near an appropriate low energy absorption maximum of said polymer for a time and at an intensity sufficient to inactivate said infectious agents.

74.A method of enhancing an image obtained from magnetic resonance imaging of a region containing a malignant tumorous growth comprising the steps of administering to a subject a non-lethal amount of a water soluble paramagnetic Mn(III) porphyrin according to claim 1, or a pharmaceutically acceptable acid form or salt thereof effective to significantly enhance the contrast in a magnetic resonance image between said malignant tumorous growth and the remainder of said image; and then imaging said region of said subject by magnetic resonance.

75. A pharmaceutical composition comprising a compound according to claim 26 and a pharmaceutically acceptable carrier or diluent.

76. A method for destroying neoplastic tissue, comprising the steps of:

contacting said tissue with a compound according to claim 1; and

subjecting said tissue to irradiation for a time and at a wavelength and intensity sufficient to destroy said tissue.

77. An electrophotographic recording material comprising a conductive support and a photosensitive layer containing at least one compound according to claim 26.

78. A thin-film device produced by electrolysis of a solution comprising a compound according to claim 1 and a surfactant for solubilizing said compound.

-54-



79. A composition comprising a photochemically oxidizable polymer containing aliphatic carbon-carbon unsaturation and an adduct or complex of a compound according to claim 1 and sulfur dioxide.

80. A composition comprising a polymeric matrix that contains at least one compound according to claim 26.


81. A method for uniform orientation of molecules in a liquid crystalline substance with respect to a substrate to which said substance is to be applied, comprising treating ~~the surface of said substrate prior to application of said~~ substance with a solution containing a compound according to claim 26.

82. An oxygen sensor comprising:
an electrically conductive substrate whose potentiometric response is substantially unaffected by the presence of oxygen gas; and
an electrolytic oxidative polymeric membrane adherently and substantially uniformly coating a surface of said electrically conductive substrate, said electrolytic oxidative polymeric membrane being formed by electrolytic polymerization and comprising at least one compound according to claim 26.


1/3

**Fig. 1****Fig. 2**

2/3

***Fig. 3***

3/3

**Fig. 4**

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US93/07686

A. CLASSIFICATION OF SUBJECT MATTER

IPC(5) :C09B 47/00; A61K 31/40

US CL :540/145; 514/185; 526/238.1, 238.2, 259; 424/9

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 252/299.61; 540/145; 514/185; 526/238.1, 238.2, 259; 424/9

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US, A, 4,988,808 (Morgan et al) 29 January 1991, see entire document.	26-35
Y, E	US, A, 5,252,730 (Mackey) 12 October 1993, see entire document.	25; 49-53, 69, 80
Y	Journal of Organic Chemistry, Vol. 54, issued 10 November 1989, "New Minnetian et al., Syntheses and Reactions of Some Halogenated Porphyrins," pages 5567-5574, see entire document.	1-24
A, P	US, A, 5,200,481 (Sounik et al) 06 April 1993, see claims.	25, 49-53, 69, 80

☒ Further documents are listed in the continuation of Box C. ☐ See patent family annex.

* Special categories of cited documents:	T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principles or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	X	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	Y	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	A	document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means		
P document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

31 OCTOBER 1993

Date of mailing of the international search report

18 NOV 1993

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. NOT APPLICABLE

Authorized officer

P.K. SRIPADA JD

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/07686

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P	US, A, 5,171,741 (Dougherty) 15 December 1992, see the abstract.	74, 76, 78
X, P	US, A, 5,164,944 (Nelson et al) 08 December 1992, see the abstract.	74, 76, 78

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/07686

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:
(Telephone Practice)
Please See Extra Sheet.

1. ☒ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US93/07686

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

- I. Claims 1-24, 26-48, 54-68, 70, 71, 75 and 79, drawn to porphyrins, process and composition, Class 540, subclass 145.
- II. Claims 25, 49-53, 69 and 80, drawn to polymers, Class 526, subclasses 238.2, 238.2, 259 and 241.
- III. Claim 72 and 73, drawn to method of inactivating infectious agents. Class 514, subclass 185.
- IV. Claims 74 and 76, method of imaging and destroying classes 424 and 514, subclasses 9 and 185.
- V. Claim 77, drawn to use as recording material, Class 430, subclass 270.
- VI. Claim 78, drawn to use as thin-film device by electrolysis, Class 430, subclass 52.
- VII. Claims 81 and 82, drawn to liquid crystals, Class 252, subclass 299.61.

These inventions lack unity because Group I is drawn to monomers and Group II is drawn to polymers which are high molecular weight compounds. Group III is drawn to inactivating infectious agents. Group IV is drawn to imaging and destroying use, namely recording material. In contrast Group VI is drawn to a thin film device produced by electrolysis. Group VII is drawn to another use as liquid crystals as sensors. Thus each Group III, IV, V, VI and VII is a different method of use.

The Patent Bank Cover Sheet

Date: October 20,1997 Time:
Customer ID:
Order Contact:
Contact Phone:
Account Bal:

Customer Name:
Customer Addr1:
Customer Addr2:
Customer City:
Customer State:
Customer Zip:

Invoice No:
Docket:
Country:
Document No:
Kind:
Total Pages:
File Wrapper:

Delivery Option:
Delivery Shipper:
Shipper Acct No: